



# Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration

ISSN: 2167-8421 (Print) 2167-9223 (Online) Journal homepage: <https://www.tandfonline.com/loi/iafd20>

## ALSUntangled 44: curcumin

Richard Bedlack & ALSUntangled Group

To cite this article: Richard Bedlack & ALSUntangled Group (2018) ALSUntangled 44: curcumin, Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration, 19:7-8, 623-629, DOI: [10.1080/21678421.2018.1440738](https://doi.org/10.1080/21678421.2018.1440738)

To link to this article: <https://doi.org/10.1080/21678421.2018.1440738>

 Published online: 01 Mar 2018.

---

 Submit your article to this journal [↗](#)

---

 Article views: 12293

---

 View related articles [↗](#)

---

 View Crossmark data [↗](#)

---

 Citing articles: 1 View citing articles [↗](#)

---



## ARTICLE

**ALSUntangled 44: curcumin**

## ALSUNTANGLED GROUP

ALSUntangled reviews alternative therapies are presented on behalf of persons with ALS (PALS). Here we review the use of curcumin for ALS, for which we have had over 100 requests (1). Curcumin is a chemical compound in the spice turmeric, accounting for ~0.5–5% of the dry weight of turmeric powder (2–4). It is also found in the curry powder used in Middle Eastern and Asian cuisine; however, the use of other spices in curry powder dilutes curcumin to significantly lower levels than in pure turmeric powder (3). Because curcumin is a small component of turmeric and an even smaller component of curry powder (3,4), we focus our review on curcumin dietary supplements.

**Overview**

Curcumin comprises the majority of several structurally similar compounds in turmeric called curcuminoids, which belong to a chemical class called polyphenols (2). Curcumin is most commonly administered orally. The medicinal utility of oral curcumin has been questioned, because it is poorly absorbed from the gut, rapidly metabolized, has been reported to inhibit several proteins important for normal physiology, and is toxic to several healthy cell lines (5,6). Different curcumin formulations have been created to try and optimize perceived problems with absorption and/or distribution within the body; these will be discussed in the “Dosing” section below. It has also been suggested that orally administered curcumin might act primarily by altering the fecal microbiome (7), which would be possibly independent of absorption. Curcumin is reported to have multiple downstream biological effects (6–10) and as a result has been tested in animal models and/or clinical trials for many different diseases, including Alzheimer’s disease, arthritis, cancer, hyperlipidemia, Parkinson’s disease, Charcot-Marie-Tooth disease, spinal cord injury, and stroke. *In vitro*

studies on curcumin have been criticized, because the compound itself is unstable in common *in vitro* testing conditions and could potentially interfere with a variety of *in vitro* assays through several mechanisms, potentially leading to false interpretations due to real bioactivity of degradation products or assay interference (5). Nonetheless, many published clinical trials with curcumin in varying formulations show clinical benefits in various diseases, suggesting that curcumin warrants further study (11).

**Mechanisms**

At least four of curcumin’s purported mechanisms of action might be relevant in treating ALS: modulation of neuroinflammation, reduction of oxidative stress, amelioration of protein aggregation, and alteration of the fecal microbiome.

*Neuroinflammation*

Neuroinflammation is thought to have a role in ALS pathogenesis, possibly through activation of the nuclear factor- $\kappa$ B (NF- $\kappa$ B) pro-inflammatory cellular pathway (12). Patients with sporadic ALS have elevated NF- $\kappa$ B expression in microglia (13). Reduced expression of NF- $\kappa$ B in microglia slows progression and extends survival in multiple mouse models of ALS (14–16). Curcumin can reportedly decrease NF- $\kappa$ B pathway activation in rat and mouse microglial cell cultures (17–19) and in rat brain when delivered to the animals via intraperitoneal injection (20) or orally (21). Curcumin also reportedly decreases NF- $\kappa$ B activation in human cell lines (22–25). Despite the promising preclinical data, a clinical trial, with cancer patients, utilizing a high bioavailability oral form of curcumin (Theracurmin) could not find a change in NF- $\kappa$ B levels in peripheral blood monocytes (26).

*Oxidative stress*

Oxidative stress may also be a key player in ALS pathology (27). PALS have elevated oxidative species and decreased antioxidants in their blood (28). Increased oxidative stress in the motor cortex correlates with worse ALS disease severity (29). Compounds that induce Nrf2-ARE, a molecular antioxidant pathway, delay disease progression in a SOD1 mutant mouse model of ALS (30–32). Curcumin reportedly neutralizes free radicals *in vitro* (33) and activates the Nrf2-ARE pathway in the brain of rats (34–36), skeletal muscle of mice (37) and rats (38), and isolated rat astrocytes (39). A single very high oral dose of curcumin (Theracurmin formulation) was able to partially prevent acute oxidative stress induced damage to the brain of mice (40). In multiple human trials, oral curcumin treatment was associated with biomarkers of increased antioxidative effect in the blood (41–51) including Nrf2 activation (51).

*Protein aggregation*

Misfolded proteins (“aggregates”), such as SOD1, are present in the motor neurons of PALS with familial and sporadic forms of the disease and some believe these play a pathogenic role (52). Treatments targeting protein aggregates can delay progression in ALS animal models (52,53). Curcumin can reportedly inhibit SOD1 aggregation *in vitro* (54), however, the concentrations of curcumin used in this experiment are so high that these concentrations would be virtually impossible to obtain in humans. Curcumin can also promote the expression of genes related to the clearance of protein aggregates in isolated blood cells from PALS and people with Alzheimer’s disease (55). Unlike neuroinflammation and oxidative stress, we found no human trials testing curcumin’s ability to affect protein aggregation.

*Alteration in fecal microbiome*

An altered fecal microbiome acting directly through a bacterial toxin or via a secondary effect on neuroinflammation through the “gut brain axis” was first postulated as a cause of ALS more than a decade ago (56). Two recent studies (57,58), but not another (59), found an altered microbiome in PALS. Altering the fecal microbiome in the mutant SOD1 mouse model of ALS prolongs survival (60) and in a worm model of ALS improves motor function (61). Oral curcumin (7) and oral Theracurmin (62) can alter the fecal microbiome in mice though it is not yet clear it can do this in a way that would help PALS.

The above-described mechanistic data are potentially exciting, but some experiments may need to be replicated given curcumin’s ability to interfere with *in vitro* assays (5). Despite these

limitations, the data are at least theoretically compelling, so ALSUntangled assigns a TOE “Mechanism” grade of C (Table 1).

**Pre-clinical models**

We found three studies, at a single institution, exploring the effect of chemical analogs of curcumin in a ALS cell model involving NSC-34 mouse spinal cord-neuroblastoma cells transfected with mutant TDP-43 or a fragment of wildtype TDP-43. In this model, either mutant TDP-43 or a fragment resulted in hyperexcitability, altered mitochondrial morphology and complex I activity, protein aggregation, and elevated a marker of oxidative stress (63–65). These studies show that certain chemical analogs of curcumin can reduce hyperexcitability (63), rescue mitochondria morphology and complex I activity (64), reduce a marker of oxidative stress and decrease protein aggregation (65). None of these studies have been independently replicated and not all experiments were adequately controlled given curcumin’s ability to interfere with *in vitro* assays (5). It is unclear how the chemical analogs that were used would perform in human trials or whether any experimental results obtained in this non-human cellular model will translate into a treatment for PALS.

Based on these three flawed published studies showing benefits of curcumin chemical analogs in an ALS cell model, ALSUntangled assigns a TOE “Pre-Clinical Models” grade of C (Table 1).

**Cases**

In the online community PatientsLikeMe, 28 PALS report taking curcumin at doses ranging from 500–1500 mg daily, and eight of these completed detailed evaluations. Only one perceived “slight” effectiveness for their ALS; the others perceived no effectiveness or could not tell that it was helping (66).

We are aware of three other PALS who experienced substantial and sustained functional improvement on cocktails of treatments that included curcumin at some point. Members of the ALSUntangled team obtained their medical records and independently validated their diagnoses and motor improvements (67). There are multiple possible explanations for their improvements, including one of the other supplements or drugs they took, endogenous resistance to ALS, or an unrecognized ALS mimic syndrome (67).

Based on the three PALS with validated diagnoses who experienced validated improvements on regimens containing curcumin, ALSUntangled assigns a TOE “Cases” grade of B (Table 1).

Table 1. Table of evidence.

	Grade	Explanation
Mechanism	C	Curcumin can theoretically act on at least four plausible ALS mechanisms.
Pre-Clinical	C	In three flawed studies using a TDP43 transfected mouse cell model of ALS, chemical analogs of curcumin have been shown to be beneficial.
Cases	B	Three PALS with validated diagnoses experienced substantial and sustained functional improvements on different multi-compound regimens that all included a form of curcumin.
Trials	C	One small published randomized, double blinded, placebo-controlled pilot trial showed some benefit with a nanoparticle formulation of curcumin.
Risks	B (oral)/D (intravenous)	Oral curcumin is associated with only rare, minor side effects. Intravenous curcumin is associated with severe adverse events including death.

## Trials

There has been one published pilot trial of curcumin in PALS which used a proprietary formulation of curcumin called Sina Curcumin (68). This was a 12-month double blinded randomized placebo-controlled trial that enrolled 54 PALS. The Sina Curcumin group took one capsule by mouth per day that contained 80 mg curcuminoids. All patients, regardless of the assigned group, took riluzole and an antacid. The results showed that the Sina Curcumin group was less likely to experience a major event, defined as death or mechanical ventilation dependency, even when controlling for other factors such as age. Despite the difference in major events, there was no significant difference found between groups in progression of ALSFRS-R score or tests of arm and leg muscle strength (MMT). The results of this study are difficult to interpret because of significant differences in age and motor testing between the placebo and Sina Curcumin groups at baseline.

We also found an unpublished pilot trial of a supplement called Brainoil (69). Brainoil is said to be a blend of curcumin, piperine, and several herbs which is processed and combined with oil into a gel. This was a six-month trial that was initially double-blind and placebo controlled and enrolled 42 PALS. After the first three months, the placebo group was crossed over to Brainoil and the trial was open-label for the remaining three months. The results showed a trend toward slower progression on ALSFRS-R scores from 0 to 6 months in the Brainoil group compared with the placebo group, but these did not reach significance. The trial was likely underpowered to detect significant changes on this scale and additionally may have suffered from a placebo effect given that it was open-label for three months, so in our opinion, the results are inconclusive.

Based on the above trials, ALSUntangled assigns a TOE “Trials” grade of C (Table 1).

## Risks

Despite the *in vitro* data suggestive that curcumin could have toxic functions (5,6), more than 120 human clinical trials have been conducted with

orally administered curcumin at doses of up to 12 g per day. We found no serious adverse events attributed to curcumin in these trials. Meta-analyses suggest that side effects of any kind are rare, affecting less than 10% of patients participating in oral curcumin trials; the most common reported side effects include abdominal pain, nausea, indigestion, constipation, diarrhea, and hot flashes (5, 70–72). Intravenous curcumin, which is far less commonly studied, is associated with at least one severe allergic reaction and one death (73). Since curcumin can affect enzymes involved in drug metabolism (5,74), it may interact with other drugs or supplements (74–77).

Based on this information, ALSUntangled assigns a TOE “Risks” grade of B for oral curcumin and D for intravenous curcumin (Table 1). It is important to note that there are many ways of formulating oral or intravenous curcumin and that these different forms could have different safety profiles. Also, while curcumin has been tested for safety in the general population, there is minimal specific experience with it in PALS.

## Dosing and costs

Although oral curcumin studies have utilized a wide range of different dosages, most were unable to detect curcumin in the serum of participants (5). Different oral formulations have been developed to try and improve blood concentrations of curcumin. The two most common strategies tested in clinical trials are creation of water soluble nanoparticles and the addition of compounds (“bioenhancers”) that improve curcumin bioavailability (78); some of these formulations with references for pharmacokinetic data are listed in Table 2. However, in light of the fact that curcumin may not need to be absorbed from the gut to work (7), it is difficult to select an optimal product for future ALS trials based on absorption data. Besides the above described trial in PALS utilizing Sina Curcumin (68), we only found one other successful trial in neurological diseases. In a trial on mild cognitive impairment (MCI), 21 patients with MCI were randomized to receive “Theracurmin” at 90 mg twice daily had significantly improved blinded measures of memory and

Table 2. Selected curcumin formulations purported to increase serum concentrations.

Product category	Product name	Description	Supporting references
Nanoparticle	Sina Curcumin	Curcuminoids in nanomicelles	No pharmacokinetic data found.
Nanoparticle	Theracurmin	Curcumin dispersed with colloidal sub-micron particles	(79,80)
Nanoparticle	Meriva	Curcumin complexed with phospholipids	(80,81)
Addition of Bioenhancer	Bio-Curmin (BCM-95)	Curcuminoids with essential oils	(80,82)
Addition of Bioenhancer	BioPerine	Curcumin with piperine (a component of black pepper)	(83)

attention over 18 months, while 19 patients randomized to placebo worsened (84). This clinical study, along with the above mentioned study of Theracurmin partially preventing acute oxidative stress induced damage in the brain of mice (40), suggests that oral Theracurmin would be a promising formulation for a future ALS trial.

The cost of curcumin varies somewhat depending on the dose and specific formulation taken. One month's supply of Theracurmin at 90 mg twice daily currently costs \$53 (85).

## Conclusion

Oral curcumin is safe, inexpensive, and has at least four potential mechanisms by which it might theoretically be useful in treating PALS. Flawed preclinical studies showed benefits of a curcumin chemical analog in a cell model of ALS, three PALS experienced validated motor improvements on regimens including curcumin (although there are several alternative explanations for these improvements) and there is one small pilot trial showing some benefit of curcumin in PALS. Based on the evidence presented in this review, some of us are planning a trial of Theracurmin at 90 mg twice daily in PALS.

## Acknowledgements

ALSUntangled Reviewers who contributed to this paper include the following: Jesse Crayle (who wrote the first draft), Richard Bedlack, Paul Barkhaus, Michael S Bereman, Greg Carter, Keelie Denson, Laurie Gutmann, Daniel Harrison, Pamela Kittrell, Christopher J McDermott, Kathy Mitchell, Sabrina Paganoni, Meraida Polak, Kristiana Salmon, and Paul Wicks.

## Declaration of interest

ALSUntangled is sponsored by the ALS Association and the Motor Neurone Disease Association. Richard Bedlack has research support from ALSA, MNDA, Cytokinetics, Neuraltus and GSK, and consulting support from ALSA, Avanir, Neuraltus, Ultragenyx, Cytokinetics, Mallinkrodt, and Brainstorm Cell. Paul Wicks (PW) is an employee of PatientsLikeMe and holds stock options in

the company. PW is an associate editor at the Journal of Medical Internet Research and is on the Editorial Boards of The BMJ and BMC Medicine. The PatientsLikeMe Research Team has received research funding (including conference support and consulting fees) from Abbvie, Accordia, Actelion, Alexion, Amgen, AstraZeneca, Avanir, Biogen, Boehringer Ingelheim, Celgene, EMD, Genentech, Genzyme, Janssen, Johnson & Johnson, Merck, Neuraltus, Novartis, Otsuka, Permobil, Pfizer, Sanofi, Shire, Takeda, Teva, and UCB. The PatientsLikeMe R&D team has received research grant funding from Kaiser Permanente, the Robert Wood Johnson Foundation, Sage Bionetworks, The AKU Society, and the University of Maryland. PW has received speaker fees from Bayer and honoraria from Roche, ARISLA, AMIA, IMI, PSI, and the BMJ.

## References

1. Available at: <http://www.alsuntangled.com/open.php>. Accessed January 1, 2018. Archived by WebCite® at: <http://www.webcitation.org/6w9UBLg03>
2. Available at: <https://en.wikipedia.org/wiki/Curcumin>. Accessed January 14, 2018. Archived by WebCite® at: <http://www.webcitation.org/6wTGY73ki>
3. Tayyem R, Heath D, Al-Delaimy W, Rock C. Curcumin content of turmeric and curry powders. *Nutr Cancer*. 2006;55:126–31.
4. Jayaprakasha G, Mohan Rao L, Sakariah K. Improved HPLC method for the determination of curcumin, demethoxycurcumin, and bisdemethoxycurcumin. *J Agric Food Chem*. 2002;50:3668–72.
5. Nelson K, Dahlin J, Bisson J, Graham J, Pauli G, Walters M. The essential medicinal chemistry of curcumin. *J Med Chem*. 2017;60:1620–37.
6. Shehzad A, Qureshi M, Anwar M, Lee Y. Multifunctional curcumin mediate multitherapeutic effects. *J Food Sci*. 2017;82:2006–15.
7. Shen L, Liu L, Hong-Fang J. Regulative effects of curcumin spice administration on gut microbiota and its pharmacological implications. *Food Nutr Res*. 2017;6:1361780.
8. Zhou H, Beevers C, Huang S. The targets of curcumin. *Curr Drug Targets*. 2011;12:332–47.
9. Amairaj A, Pius A, Gopi S, Gopi S. Biological activities of curcuminoids, other biomolecules from tumeric and their derivatives: a review. *J Tradit Complement Med*. 2016; 7:205–33.
10. Patzkó A, Bai Y, Saporta MA, Katona I, Wu X, Vizzuso D, et al. Curcumin derivatives promote Schwann cell differentiation and improve neuropathy in R98C CMT1B mice. *Brain*. 2012;135:3551–66.

11. Heger M. Drug screening: Don't discount all curcumin trial data. *Nature*. 2017;543:40.
12. Hovden H, Frederiksen J, Pedersen S. Immune system alterations in amyotrophic lateral sclerosis. *Acta Neurologica Scandinavica*. 2013;128:287–96.
13. Sako W, Ito H, Yoshida M, Koizumi H, Kamada M, Fujita K, et al. Nuclear factor  $\kappa$ B expression in patients with sporadic amyotrophic lateral sclerosis and hereditary amyotrophic lateral sclerosis with optineurin mutations. *Clin Neuropathol*. 2012;31:418–23.
14. Frakes A, Ferraiuolo L, Haidet-Phillips A, Schmelzer L, Braun L, Miranda C, et al. Microglia induce motor neuron death via the classical NF- $\kappa$ B pathway in amyotrophic lateral sclerosis. *Neuron*. 2014;81:1009–23.
15. Patel P, Julien J, Kriz J. Early-stage treatment with withaferin a reduces levels of misfolded superoxide dismutase 1 and extends lifespan in a mouse model of amyotrophic lateral sclerosis. *Neurotherapeutics*. 2015;12:217–33.
16. Swarup V, Phaneuf D, Dupre N, Petri S, Strong N, Kriz J, et al. Deregulation of TDP-43 in amyotrophic lateral sclerosis triggers nuclear factor  $\kappa$ B-mediated pathogenic pathways. *J Exp Med*. 2011;208:2429–47.
17. Jin C, Lee J, Park C, Choi Y, Kim G. Curcumin attenuates the release of pro-inflammatory cytokines in lipopolysaccharide-stimulated BV2 microglia. *Acta Pharmacologica Sinica*. 2007;28:1645–51.
18. Kang G, Kong P, Yuh Y, Lim S, Yim S, Chun W, et al. Curcumin suppresses lipopolysaccharide-induced cyclooxygenase-2 expression by inhibiting activator protein 1 and nuclear factor B bindings in BV2 microglial cells. *J Pharmacol Sci*. 2004;94:325–8.
19. Yang S, Zhang D, Yang Z, Hu X, Qian S, et al. Curcumin protects dopaminergic neuron against LPS induced neurotoxicity in primary rat neuron/glia culture. *Neurochem Res*. 2008;33:2044–53.
20. Sharma N, Sharma S, Nehru B. Curcumin protects dopaminergic neurons against inflammation-mediated damage and improves motor dysfunction induced by single intranigral lipopolysaccharide injection. *Inflammopharmacol*. 2017;25:351–68.
21. Tiwari V, Chopra K. Protective effect of curcumin against chronic alcohol-induced cognitive deficits and neuroinflammation in the adult rat brain. *Neuroscience*. 2013;244:147–58.
22. Reuter S, Charlet J, Juncker T, Teiten M, Dicato M, Diederich M. Effect of curcumin on nuclear factor kappaB signaling pathways in human chronic myelogenous K562 leukemia cells. *Ann N Y Acad Sci*. 2009;1171:436–47.
23. Jobin C, Bradham C, Russo M, Juma B, Narula A, Brenner D, et al. Curcumin blocks cytokine-mediated NF- $\kappa$ B activation and proinflammatory gene expression by inhibiting inhibitory factor I- $\kappa$ B kinase activity. *J Immunol*. 1999;163:3474–83.
24. Schwertheim S, Wein F, Lennartz K, Worm K, Schmid K, Sheu-Grabellus Y. Curcumin induces G2/M arrest, apoptosis, NF- $\kappa$ B inhibition, and expression of differentiation genes in thyroid carcinoma cells. *J Cancer Res Clin Oncol*. 2017;143:1143–54.
25. Pimentel-Gutiérrez H, Bobadilla-Morales L, Barba-Barba C, Ortega-De-La-Torre C, Sánchez-Zubieta F, Corona-Rivera J, et al. Curcumin potentiates the effect of chemotherapy against acute lymphoblastic leukemia cells via downregulation of NF- $\kappa$ B. *Oncol Lett*. 2016;12:4117–24.
26. Kanai M, Otsuka Y, Otsuka K, Sato M, Nishimura T, Mori Y, et al. A phase I study investigating the safety and pharmacokinetics of highly bioavailable curcumin (Theracurmin) in cancer patients. *Cancer Chemother Pharmacol*. 2013;71:1521–30.
27. Bozzo F, Mirra A, Carri M. Oxidative stress and mitochondrial damage in the pathogenesis of ALS: new perspectives. *Neurosci Lett*. 2017;636:3–8.
28. Blasco H, Garçon G, Patin F, Veyrat-Durebex C, Boyer J, Devos D, et al. Panel of oxidative stress and inflammatory biomarkers in ALS: a pilot study. *Can J Neurol Sci*. 2017;44:90–5.
29. Ikawa M, Okazawa H, Tsujikawa T, Matsunaga A, Yamamura O, Mori T, et al. Increased oxidative stress is related to disease severity in the ALS motor cortex. *Neurology*. 2015;84:2033–9.
30. Neymotin A, Calingasan N, Wille E, Naseri N, Petri S, Damiano M, et al. Neuroprotective effect of Nrf2/ARE activators, CDDO ethylamide and CDDO trifluoroethylamide, in a mouse model of amyotrophic lateral sclerosis. *Free Rad Biol Med*. 2011;51:88–96.
31. Mead R, Higginbottom A, Allen S, Kirby J, Bennet E, Barber S, et al. S[+] Apomorphine is a CNS penetrating activator of the Nrf2-ARE pathway with activity in mouse and patient fibroblast models of amyotrophic lateral sclerosis. *Free Rad Biol Med*. 2013;61:438–52.
32. Tanaka K, Kanno T, Yanagisawa Y, Yasutake K, Inoue S, Hirayama N, et al. A novel acylaminoimidazole derivative, WN1316, alleviates disease progression via suppression of glial inflammation in ALS mouse model. *PLoS One*. 2014;9:e87728.
33. Ak T, Gulcin A. Antioxidant and radical scavenging properties of curcumin. *Chem Biol Interact*. 2008;174:27–37.
34. Cui Q, Li X, Zhu H. Curcumin ameliorates dopaminergic neuronal oxidative damage via activation of the Akt/Nrf2 pathway. *Mol Med Rep*. 2016;13:1381.
35. Li W, Suwanwela N, Patumraj S. Curcumin by down-regulating NF- $\kappa$ B and elevating Nrf2, reduces brain edema and neurological dysfunction after cerebral I/R. *Microvasc Res*. 2016;106:117–27.
36. Wu J, Li Q, Wang X, Yu S, Li L, Wu X, et al. Neuroprotection by curcumin in ischemic brain injury involves the Akt/Nrf2 pathway. *PLoS One*. 2013;8:e59843.
37. He H, Wang G, Gao Y, Ling W, Yu Z, Jin T. Curcumin attenuates Nrf2 signaling defect, oxidative stress in muscle and glucose intolerance in high fat diet-fed mice. *World J Diabetes*. 2012;3:94–104.
38. Sahin K, Pala R, Tuzcu M, Ozdemir O, Orhan C, Sahin N, et al. Curcumin prevents muscle damage by regulating NF- $\kappa$ B and Nrf2 pathways and improves performance: an in vivo model. *J Inflamm Res*. 2016;9:147–54.
39. Jiang H, Tian X, Guo Y, Duan W, Bu H, Li C. Activation of nuclear factor erythroid 2-related cytoprotective signaling by curcumin protect primary spinal cord astrocytes against oxidative toxicity. *Biol Pharm Bull*. 2011;34:1194–7.
40. Nazari QA, Takada-Takatori Y, Hashimoto T, Imaizumi A, Izumi Y, Kume T. Potential protective effect of highly bioavailable curcumin on an oxidative stress model induced by microinjection of sodium nitroprusside in mice brain. *Food Func*. 2014;5:984–9.
41. Pungcharoenkul K, Thongnopnua P. Effect of different curcuminoid supplement dosages on total in vivo antioxidant capacity and cholesterol levels of healthy human subjects. *Phytother Res*. 2011;25:1721–6.
42. Sahebkar A, Mohammadi A, Atabati A, Rahiman S, Tavallaie S, Iranshahi M, et al. Curcuminoids modulate pro-oxidant-antioxidant balance but not the immune response to heat shock protein 27 and oxidized LDL in obese individuals. *Phytother Res*. 2013;27:1883–8.
43. Kalpravidh R, Siritanaratkul N, Insain P, Charoensakdi R, Panichkul N, Hatairaktham S, et al. Improvement in oxidative stress and antioxidant parameters in  $\beta$ -thalassaemia/Hb E patients treated with curcuminoids. *Clin Biochem*. 2010;43:424–9.
44. Takahashi M, Suzuki K, Kim H, Otsuka Y, Imaizumi A, Miyashita M, et al. Effects of curcumin supplementation on exercise-induced oxidative stress in humans. *Int J Sports Med*. 2014;35:469–75.

45. Usharani P, Mateen A, Naidu M, Raju Y, Chandra N. Effect of NCB-02, atorvastatin, and placebo on endothelial function, oxidative stress, and inflammatory markers in patients with type 2 diabetes mellitus. *Drugs RD*. 2008;9:243–50.
46. Panahi P, Ghanei M, Hajhashemi A, Sahebkar A. Effects of curcuminoids-piperine combination on systemic oxidative stress, clinical symptoms and quality of life in subjects with chronic pulmonary complications due to sulfur mustard: a randomized controlled trial. *J Diet Suppl*. 2016;13:93–105.
47. Panahi Y, Alishiri G, Parvin S, Sahebkar A. Mitigation of systemic oxidative stress by curcuminoids in osteoarthritis: results of a randomized controlled trial. *J Diet Suppl*. 2016;13:209–20.
48. Panahi Y, Khalili N, Sahebi E, Namazi S, Karimian M, Majeed M, et al. Antioxidant effects of curcuminoids in patients with type 2 diabetes mellitus: a randomized controlled trial. *Inflammopharmacol*. 2017;25:25–31.
49. Panahi Y, Hosseini M, Khalili N, Naimi E, Majeed M, Sahebkar A. Antioxidant and anti-inflammatory effects of curcuminoid-piperine combination in subjects with metabolic syndrome: a randomized controlled trial and an updated meta-analysis. *Clin Nutr*. 2015;34:1101–8.
50. Hejazia J, Rastmaneshb R, Taleban FA, Molana SH, Hejazi E, Ehtejab G, et al. Effect of curcumin supplementation during radiotherapy on oxidative status of patients with prostate cancer: a double blinded, randomized, placebo-controlled study. *Nutr Cancer*. 2016;68:77–85.
51. Yang H, Xu W, Zhou Z, Liu J, Li X, Chen L, et al. Curcumin attenuates urinary excretion of albumin in type II diabetic patients with enhancing nuclear factor erythroid-derived 2-like 2 (Nrf2) system and repressing inflammatory signaling efficacies. *Exp Clin Endocrinol Diabetes*. 2015;123:360–7.
52. Parakh S, Atkin J. Protein folding alterations in amyotrophic lateral sclerosis. *Brain Res*. 2016;1648:633–49.
53. Kieran D, Kalmar B, Dick J, Riddoch-Contreras J, Burnstock G, Greensmith L. Treatment with arimoclomol, a coinducer of heat shock proteins, delays progression in ALS mice. *Nat Med*. 2004;10:402–5.
54. Curcumin binds to the pre-fibrillar aggregates of Cu/Zn superoxide dismutase (SOD1) and alters its amyloidogenic pathway resulting in reduced cytotoxicity. *Biochim Biophys Acta*. 2015;1854:426–36.
55. Cashman J, Gagliardi S, Lanier M, Ghirmai S, Abel K, Fiala M. Curcumins promote monocytic gene expression related to beta-amyloid and superoxide dismutase clearance. *Neurodegenerative Dis*. 2012;10:274–6.
56. Longstreth J, Meschke J, Davidson S, Smoot L, Koepsell T. Hypothesis: a motor toxin produced by a clostridial species residing the gut causes ALS. *Med Hypotheses*. 2005;64:1153.
57. Rowin J, Xia Y, Sun J. Gut inflammation and dysbiosis in human motor neuron disease. *Physiol Rep*. 2017;5:e13443.
58. Mazzini L, Mogna L, Marchi FD, Amoroso A, Pane M, Aloiso I, et al. Potential role of gut microbiota in ALS pathogenesis. *Amyotroph Lateral Scler and Frontotemporal Degenr*. 2017;18:245.
59. Brenner D, Hiergeist A, Adis C, Mayer B, Gessner A, Ludolph A, et al. The fecal microbiome of ALS patients. *Neurobiol Aging*. 2018;61:132–7.
60. Zhang Y, Wu S, Yi J, Xia Y, Jin D, Zhou J, et al. Target intestinal microbiota to alleviate disease progression in amyotrophic lateral sclerosis. *Clin Ther*. 2017;39:322–36.
61. Labarre A, Guitard E, Parker J. Variation of gut microbiome rescues paralysis and neurodegeneration profiles in *C. elegans* ALS models. *Amyotroph Lateral Scler and Frontotemporal Degenr*. 2017;18:245–6.
62. Ohno M, Nishida A, Sugitani Y, Nishino K, Inatomi O, Sugimoto M et al. Nanoparticle curcumin ameliorates experimental colitis via modulation of gut microbiota and induction of regulatory T cells. *PLoS One*. 2017;12:e0185999.
63. Dong H, Xu L, Wu L, Wang X, Duan W, Li H, et al. Curcumin abolishes mutant TDP-43 induced excitability in a motoneuron-like cellular model of ALS. *Neuroscience*. 2014;272:141–53.
64. Lu J, Duan W, Guo Y, Jiang H, Li Z, Huang J, et al. Mitochondrial dysfunction in human TDP-43 transfected NSC34 cell lines and the protective effect of dimethoxy curcumin. *Brain Res Bull*. 2012;89:185–90.
65. Duan W, Guo Y, Xiao J, Chen X, Li Z, Han H, et al. Neuroprotection by monocarbonyl dimethoxycurcumin C: ameliorating the toxicity of mutant TDP-43 via HO-1. *Mol Neurobiol*. 2014;49:368–79.
66. Available at: <https://www.patientslikeme.com/treatments/show/532>. Accessed: January 8, 2018. Archived by WebCite® at: <http://www.webcitation.org/6wJwJcnpN>
67. Harrison D, Bedlack R. ALS reversals: demographics, disease characteristics, treatments and co-morbidities. Presented at American Academy of Neurology April 25, 2017.
68. Ahmadi M, Elmira A, Nafissi S, Jaafari MR, Harirchian MH, Sarraf P, et al. Safety and efficacy of nanocurcumin as add-on therapy to riluzole in patients with amyotrophic lateral sclerosis: a pilot randomized clinical trial. *Neurotherapeutics*. 2018. [Epub ahead of print]. DOI: 10.1007/s13311-018-0606-7
69. Lo Gerfo A, Caldarazzo I, Petrozzi L, Rocchi A, Modena M, Pasquinelli A, et al. Amyotrophic lateral sclerosis and oxidative stress: a double blind trial after dietary supplement by a curcumin donor. Presented at 2nd Congress of European Academy of Neurology May 28–31, 2016.
70. Qin S, Huang L, Gong J, Shen S, Huang J, Ren H, et al. Efficacy and safety of turmeric and curcumin in lowering blood lipid levels in patients with cardiovascular risk factors: a meta-analysis of randomized controlled trials. *Nutr J*. 2017;16:68.
71. Sahebkar A, Henrotin Y. Analgesic efficacy and safety of curcuminoids in clinical practice: a systematic review and meta-analysis of randomized controlled trials. *Pain Med*. 2016;17:1192–202.
72. Al-Karawi D, Al Mamoori D, Tayyar Y. The role of curcumin administration in patients with major depressive disorder: mini meta-analysis of clinical trials. *Phytother Res*. 2016;30:175–83.
73. Available at: <https://www.fda.gov/Drugs/Guidance/ComplianceRegulatoryInformation/PharmacyCompounding/ucm570192.htm>. Accessed: January 14, 2018. Archived by WebCite® at: <http://www.webcitation.org/6wTFyQ5bh>
74. Bahramsoltani R, Rahimi R, Farzaei MH. Pharmacokinetic interactions of curcuminoids with conventional drugs: a review. *J Ethnopharmacol*. 2017;209:1–12.
75. Kusuvara H, Furuie H, Inano A, Sunagawa A, Yamada S, Wu C, et al. Pharmacokinetic interaction study of sulphasalazine in healthy subjects and the impact of curcumin as an in vivo inhibitor of BCRP. *Brit J Pharmacol*. 2012;166:1793–803.
76. Chen Y, Liu WH, Chen BL, Fan L, Han Y, Wang G, et al. Plant polyphenol curcumin significantly affects CYP1A2 and CYP2A6 activity in healthy, male chinese volunteers. *Ann Pharmacother*. 2010;44:1038–45.
77. Juan H, Terhaag B, Cong Z, Bi-Kui Z, Rong-Hua Z, et al. Unexpected effect of concomitantly administered curcumin on the pharmacokinetics of talinolol in healthy Chinese volunteers. *Eur J Clin Pharmacol*. 2007;63:663–8.
78. Mahran RI, Hagraas MM, Sun D, Brenner DE. Bringing curcumin to the clinic in cancer prevention: a review of strategies to enhance bioavailability and efficacy. *AAPS J*. 2017;19:54–81.

79. Kanai M, Imaizumi A, Otsuka Y, Sasaki H, Hasiguchi M, Tsujiko K, et al. Dose-escalation and pharmacokinetic study of nanoparticle curcumin, a potential anticancer agent with improved bioavailability, in healthy volunteers. *Cancer Chemother Pharmacol.* 2012;69:65–70.
80. Sunagawa Y, Hirano S, Katanasaka Y, Miyazaki Y, Funamoto M, Okamura N, et al. Colloidal submicron particle curcumin exhibits high absorption efficiency—a double blind, 3 way crossover study. *J Nutr Sci Vitaminol.* 2015;61:37–44.
81. Cuomo J, Appendino G, Dern AS, Schneider E, McKinnon TP, Brown MJ, et al. Comparative absorption of a standardized curcuminoid mixture and its lecithin formulation. *J Nat Prod.* 2011;74:664–9.
82. Available from: <http://dolcas-biotech.com/products/bcm95/>. Accessed: January 15, 2018. Archived by WebCite® at: <http://www.webcitation.org/6wV3tc5py>
83. Shoba G, Joy D, Joseph T, Majeed M, Rajendran R, Srinivas P. Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers. *Planta Med.* 1998;64:353–6.
84. Small G, Siddarth P, Zhaoping L, Miller K, Ercoli L, Emerson N, et al. Memory and brain amyloid and tau effects of a bioavailable form of curcumin in non-demented adults: a double blind, placebo-controlled 18-month trial. *Am J Geriatr Psychiatry.* 2017;17:S1064.
85. Available at: <http://www.integrativepro.com/Products/Metabolic/Theracurmin-HP>. Accessed January 15, 2018. Archived by WebCite® at: <http://www.webcitation.org/6wUXG11HH>